



A Commercial-Stage Pharmaceutical Company Pioneering Novel Cancer Therapies

June 2023

Forward-looking Statements and Other Important Information

This presentation contains forward-looking statements within the meaning of The Private Securities Litigation Reform Act of 1995. Such forward-looking statements include those regarding Karyopharm's guidance on its 2023 total revenue, 2023 U.S. net product revenue and 2023 non-GAAP R&D and SG&A expenses; Karyopharm's expected cash runway; the ability of selinexor or eltanexor to treat patients with multiple myeloma, endometrial cancer, myelofibrosis, myelodysplastic neoplasms, diffuse large B-cell lymphoma, and other diseases; expectations related to future clinical development and potential regulatory submissions of selinexor or eltanexor; expectations with respect to commercialization efforts; submissions to, and the review and potential approval of selinexor, eltanexor or any of its other product candidates by, regulatory authorities, including the Company's regulatory strategy, the anticipated availability of data to support such submissions, timing of such submissions and actions by regulatory authorities and the potential availability of accelerated approval pathways; the expected design of the Company's clinical trials; and the therapeutic potential of and potential clinical development plans for Karyopharm's product candidates, especially selinexor and eltanexor. Such statements are subject to numerous important factors, risks and uncertainties, many of which are beyond Karyopharm's control, that may cause actual events or results to differ materially from Karyopharm's current expectations. For example, there can be no guarantee that Karyopharm will successfully commercialize XPOVIO or that any of Karyopharm's drug candidates, including selinexor and eltanexor, will successfully complete necessary clinical development phases or that development of any of Karyopharm's drug candidates will continue. Further, there can be no guarantee that any positive developments in the development or commercialization of Karyopharm's drug candidate portfolio will result in stock price appreciation. Management's expectations and, therefore, any forward-looking statements in this press release could also be affected by risks and uncertainties relating to a number of other factors, including the following: the risk that the COVID-19 pandemic could disrupt Karyopharm's business more severely than it currently anticipates, including by negatively impacting sales of XPOVIO, interrupting or delaying research and development efforts, impacting the ability to procure sufficient supply for the development and commercialization of selinexor or other product candidates, delaying ongoing or planned clinical trials, impeding the execution of business plans, planned regulatory milestones and timelines, or inconveniencing patients; the adoption of XPOVIO in the commercial marketplace, the timing and costs involved in commercializing XPOVIO or any of Karyopharm's drug candidates that receive regulatory approval; the ability to obtain and retain regulatory approval of XPOVIO or any of Karyopharm's drug candidates that receive regulatory approval; Karyopharm's results of clinical trials and preclinical studies, including subsequent analysis of existing data and new data received from ongoing and future studies; the content and timing of decisions made by the U.S. Food and Drug Administration and other regulatory authorities, investigational review boards at clinical trial sites and publication review bodies, including with respect to the need for additional clinical studies; the ability of Karyopharm or its third party collaborators or successors in interest to fully perform their respective obligations under the applicable agreement and the potential future financial implications of such agreement; Karyopharm's ability to enroll patients in its clinical trials; unplanned cash requirements and expenditures; development or regulatory approval of drug candidates by Karyopharm's competitors for products or product candidates in which Karyopharm is currently commercializing or developing; and Karyopharm's ability to obtain, maintain and enforce patent and other intellectual property protection for any of its products or product candidates. These and other risks are described under the caption "Risk Factors" in Karyopharm's Quarterly Report on Form 10-Q for the quarter ended March 31, 2023, which was filed with the Securities and Exchange Commission (SEC) on May 4, 2023, and in other filings that Karyopharm may make with the SEC in the future. Any forward-looking statements contained in this presentation speak only as of the date hereof, and, except as required by law, Karyopharm expressly disclaims any obligation to update any forward-looking statements, whether as a result of new information, future events or otherwise. Karyopharm regularly uses its website to post information regarding its business, drug development programs and governance. Karyopharm encourages investors to use www.karyopharm.com, particularly the information in the section entitled "Investors," as a source of information about Karyopharm. References to www.karyopharm.com in this presentation are not intended to, nor shall they be deemed to, incorporate information on www.karyopharm.com into this presentation by reference. Other than the currently approved indications of XPOVIO, selinexor and eltanexor are investigational drugs that have not been approved by the FDA or any other regulatory agency, and the safety and efficacy of these drugs has not been established by any agency.

XPOVIO® (selinexor) and NEXPOVIO® (selinexor) are registered trademarks of Karyopharm Therapeutics Inc. Any other trademarks referred to in this presentation are the property of their respective owners. All rights reserved.

Innovation and Patient Focused

Founded in 2008, building on over a decade of research into selective inhibition of nuclear export (SINE) as a novel mechanism of action



Passionately driven in its mission to positively impact lives and defeat cancer

Positioned for Next Stage of Growth



XPOVIO / NEXPOVIO

Approved in Multiple Myeloma (MM) and DLBCL¹

- Expanded global footprint with regulatory approvals in > 40 countries
- Total revenues expected to be \$145-160M in 2023
- Moving into earlier lines of therapy in MM

Focused Clinical Pipeline with One Planned and Two Ongoing Pivotal Studies; Optimizing Dose for Efficacy and Tolerability

- Phase 3 selinexor+ruxolitinib in treatment-naïve MF (planned)²
- Phase 3 SPd³ in R/R MM post anti-CD38
- Phase 3 selinexor as maintenance in *TP53* wild-type EC⁴

Strong Financial Position

- Cash position of \$262M⁵ at end of Q1 2023
- Cash runway until late 2025

First Quarter 2023 and Recent Highlights



**Expanding on
Multiple Myeloma
Foundation Globally**

1Q 2023 U.S. XPOVIO Net Product Revenue: \$28.3M consistent with 1Q 2022

- Continued YoY growth in total demand and community setting in a highly competitive market
- Net product revenue in 1Q 2023 adversely impacted by:
 - Increased use of the KaryForward patient assistance program (PAP)
 - Higher gross to net (GTN)

Continued Global Expansion of Selinexor

- Amended Menarini license agreement to add selected Middle East and Africa regions
- Received UK full marketing authorization for NEXPOVIO® in 2L+ MM



**Continue to Advance
Clinical Pipeline in
Core Indications**

Myelofibrosis (Selinexor)

- Strong efficacy and manageable tolerability with 60mg dose; Phase 1 data presented at AACR
 - 79% SVR35 and 58% TSS50 at Week 24 (ITT)
- Planning to initiate pivotal Phase 3 study in 1H 2023

MDS (Eltanexor)

- Encouraging interim efficacy data from Phase 2 study in hard-to-treat higher risk R/R MDS patient population
 - 8.7 months median overall survival (mOS)

Nucleus

Nuclear pore complex

XPO1

RAN-GTP

RNA binding adaptor (e.g., eIF4E)

Cytoplasm

Protein cargoes: e.g., tumor suppressor proteins and growth regulators like p53, p27, or FOXO family proteins^{1,2}

RAN-GDP (GTP hydrolysis)

RAN-GDP (GTP hydrolysis)

RNA cargoes: e.g., oncoprotein mRNA like MYC³

Exportin 1 (XPO1) transports proteins and protein-RNA complexes out of the nucleus

Adapted from Azizian NG, et al (2020)

Selinexor and Eltanexor (SINE compounds) selectively inhibit nuclear export by binding XPO1

1. Increases nuclear levels of **tumor suppressor proteins** and their activation^{4,5}
2. Traps **oncoprotein mRNA** in the nucleus, leading to reduced oncoprotein levels⁶
3. Retains **activated glucocorticoid receptor** in the nucleus, leading to altered expression of genes involved in inflammatory pathways⁷

Reduced proliferation and increased apoptosis of cancer cells⁸

SINE: Selective inhibition of nuclear export

SINE compound

Nucleus

Two Differentiated, Complementary Novel SINE Compounds



Selinexor

**First novel
XPO1 inhibitor**

- FDA-approved in MM and in DLBCL¹
- Currently being investigated in both solid tumors and hematologic malignancies²
- Penetrates blood-brain barrier (BBB)³
- Dosing: once weekly in MM (XVd), twice weekly in MM (Xd) and in DLBCL¹
- Prioritization of clinical development in *TP53* wild type endometrial cancer and myelofibrosis

Focused on areas where a **highly potent** XPO1 impact is needed



Eltanexor

**Second novel
XPO1 inhibitor**

- **Investigational compound**
- Currently being investigated in relapsed/refractory (R/R) MDS²
- Compared to selinexor, lower BBB penetration observed in select animal models^{3,†‡}
- Dosing: 5 times per week in Ph 1/2⁴
- Strong rationale for further development in both solid tumors and hematology

Focused on areas where **continuous** XPO1 inhibition needed

Prioritized and Targeted Core Indications Focused on Driving Improved Patient Outcomes in Areas of High Unmet Need

MULTIPLE MYELOMA

Enabling a **'Class Switch'** in **Earlier Lines of Therapy** to continue improving patient outcomes

ENDOMETRIAL CANCER

Potential to be the **First Maintenance-Only Treatment Option** for patients whose tumors are p53 wild-type

MYELOFIBROSIS


Potential to improve outcomes in **patients with MF**

MYELOYDYSPLASTIC NEOPLASMS


Potential to improve outcomes in **patients with MDS**

Opportunity to expand into additional lines of therapy in all four core indications

Progressing Focused Pipeline Across Cancers With High Unmet Needs



Optimizing Selinexor Dose to Improve Patient Experience and Overall Benefit



MULTIPLE MYELOMA



Selinexor Improves Progression Free Survival for Patients with Relapsed Multiple Myeloma¹

What is Multiple Myeloma?

- Cancer of the plasma cells and the **second most common blood cancer** in the world²
- Malignant plasma cells produce a paraprotein (an inactive antibody known also as M-protein) that adversely affects bone marrow, bones, and kidneys


Treatment Landscape

- Following first line progression, **treatment decisions are primarily based on physician and patient choice**, in addition to treatment guidelines
- Current standard of care is to **switch drug classes** once a regimen stops responding

Opportunity and Unmet Need


- Retrospective analyses of currently approved combination regimens demonstrate **poor outcomes for patients with multiple myeloma refractory to prior daratumumab (anti-CD38 mAb)** treatment, including low ORRs and short PFS
- **25%** patients have multiple myeloma with high-risk cytogenetics and poorer outcomes with currently available therapies⁴

Affects ~47,000 patients in the US in 2L+ ³



Striving to be a Standard of Care in 2L+, Driving Sustainable and Long-Term Growth

~47,000 US patients/year in 2L+¹



Karyopharm's focus

2

XVd

BOSTON approved
in 2L+ (Dec 2020)

3

SPd

Advance Ph 3 with all-
oral triplet targeting
2L+ and report top line
results in 2H 2024


1

Xd

STORM approved in 4L+ (July 2019)

The safety and efficacy of SPd has not been established and has not been approved by the FDA or any other regulatory authority. Combinations other than XVd and Xd are not promoted by Karyopharm but may be considered for future indication updates.

Clear Positioning within Community & Academic Settings with XPOVIO Serving an Unmet Need as a Novel Class of Therapy and Convenient Oral Treatment for 2L–4L R/R MM Post Anti-CD38 mAb



XPOVIO combinations other than XVd and Xd will not be promoted by Karyopharm, but may be considered for future indication updates.
Safety and efficacy of selinexor in combinations other than XVd and Xd have not been established and have not been approved by the US FDA or any other regulatory authority.

Generating Evidence on the Role and Effectiveness of Selinexor pre and post T-cell Mediated Therapies

Published Studies

Selinexor maintains T-cells function in mice;¹ pre-treatment can maintain effectiveness of CAR-T therapies ^{2,3}

Preclinical Research

Impact of SINE mechanism on T-cell fitness via collaborations with academic institutions

Real-World Evidence

Effectiveness of CAR-T following selinexor therapy

Clinical Research

Evaluating selinexor pre or post BCMA/CAR-T therapy

XPOVIO Update: 1Q 2023

YoY Growth in Total Demand¹ of 6%; Net Revenue Adversely Impacted by Increased Utilization of PAP and Higher GTN




1Q 2023 Highlights (YoY)

- Total demand growth¹ of 6% year over year driven by the community setting which contributed ~ 70% of revenues in Q1 2023
- Sustained business in Academics despite increased competition from T-cell therapies
- Net XPOVIO revenues adversely impacted by:
 - Significant increase in utilization of PAP (free drug) due to recent lack of funding to foundations for multiple myeloma patients
 - Higher GTN driven by increased Medicare & Medicaid rebates, and 340B discounts
- Approaching 60% of new XPOVIO patients in 2-4L²
- Improvement in tolerability perception in 2-4L per intent to prescribe data³
- Net U.S. XPOVIO revenue guidance revised to \$110-\$125M due to expected significant increase in PAP utilization
 - In 2024, significantly fewer patients expected to utilize PAP for co-pay assistance due to IRA⁴ related re-design of Part D benefits

XPOVIO® /NEXPOVIO® Approved in > 40 Countries

Country/Region	Indication(s)	Partner
Approvals		
United States		—
Europe ¹		Menarini
UK		Menarini
Mainland China		Antengene
South Korea		Antengene
Australia		Antengene
Singapore		Antengene
Canada		Forus
Israel		Neopharm
Taiwan		Antengene




-  2L+ multiple myeloma and R/R DLBCL*
-  2L+ multiple myeloma
-  Penta- or triple-class-refractory multiple myeloma and R/R DLBCL*
-  Penta- or triple-class-refractory multiple myeloma

* DLBCL approved in the U.S. under accelerated approval pathway

1. The 27 countries comprising the European Union, plus Iceland, Norway, Northern Ireland and Lichtenstein.

Phase 3 Study (XPORT-MM-031/ EMN29) Evaluating Selinexor, Pomalidomide and Dexamethasone (SPd) in Patients with Previously Treated Multiple Myeloma



Top-line data expected in 2H 2024

40mg selinexor dose was based upon evaluation of the safety and benefit of selinexor 40 and 60 mg doses in combo with Pd observed in the STOMP and 028 studies



ENDOMETRIAL CANCER

Endometrial Cancer is the Most Common Gynecologic Cancer with Significant Unmet Need for Patients with Advanced or Recurrent Disease

What is Endometrial Cancer?

- Arises from the endometrium, the layer of cells that form the lining of the uterus.

Treatment Landscape


- First-line treatment is chemotherapy (taxane plus platinum), where response rates can be as high as 67%³
- Following chemotherapy, NCCN Guidelines® recommend “watch and wait” until disease relapses⁴
- ~50% of patients with advanced or recurrent disease have *TP53* wild-type tumors⁵

Unmet Need

- Prognosis is poor, with **progression expected within ~4 months⁶** for patients responding to first-line chemotherapy treatment

The safety and efficacy of selinexor in endometrial cancer has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Nearly 66,000 new and 14,000 advanced cases diagnosed in the U.S. in 2022¹ and more than 130,000 cases in Europe²



Potential for Significant Paradigm Shift for the Treatment of Women with Advanced or Recurrent *TP53* Wild-Type Endometrial Cancer

Phase 3 SIENDO study

Generated strong hypothesis in patients with *TP53* wild-type EC

Addressing a significant unmet need


Currently no FDA approved treatments in the maintenance setting

Significant market opportunity

~14K patients diagnosed with advanced and recurrent endometrial cancer in the U.S. each year¹
~50% of these patients have *TP53* wild-type EC²


Supportive Mechanism of Action

Forced retention of p53 in the cell nucleus by inhibition of XPO1 allows p53 to carry out its tumor suppressor and other regulatory functions



The safety and efficacy of selinexor in endometrial cancer has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Updated Exploratory Subgroup Analysis from SIENDO Study in Patients with TP53 Wild Type Endometrial Cancer^{1,2} Further Supports Rationale for Evaluating Selinexor as Maintenance Therapy



Median PFS
Selinexor (n=77): **20.8 months**
Placebo (n=36): **5.2 months**

- Most common adverse events (AE) were nausea, vomiting and diarrhea with no meaningful change in the AE profile since last data cut
- Most common Gr 3/4 AEs were neutropenia (18%), nausea (12%), thrombocytopenia (9%) and fatigue (8%). No Grade 5 events observed
- AEs were generally manageable with supportive care and dose modifications


The safety and efficacy of selinexor in endometrial cancer has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

1. TP53 WT/MUT status of patients under continued evaluation. 10% of patients on the selinexor arm and 7% of patients on the control arm have no tumor TP53 status.
2. Preliminary data from pre-specified exploratory subgroup analysis as of the database cut date of November 30, 2022.

Initiated Global Phase 3 Pivotal Study; *TP53* Mutation Status Is Assessed by Companion Diagnostic Partner Foundation Medicine¹

XPORT-EC-042 Global Phase 3, Randomized, Double-Blind, Trial of Selinexor as Maintenance Therapy for Patients with *TP53* Wild-type, Advanced or Recurrent Endometrial Cancer (N=220)

Study in Collaboration with ENGOT² and GOG³



- HR-QoL, health-related quality of life; OS, overall survival; PFS, progression-free survival; PD, progressive disease; QW, every week

The safety and efficacy of selinexor in endometrial cancer has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

1. Utilizing Foundation Medicine's tissue-based next generation sequencing test to identify *TP53* status

2. European Network for Gynaecological Oncological Trial groups; 3. Gynecologic Oncology (GOG) Foundation

4. Selected secondary endpoints

MYELOFIBROSIS



XPO1, a Novel and Potentially Fundamental Mechanism in MF, has the Opportunity to Transform 1L Treatment Paradigm

Treatment Landscape and Unmet Need

~20,000 living with MF in the U.S and ~ 17,000 in EU¹

No other approved class of therapy other than JAK inhibitors; ruxolitinib generates > \$2 billion revenues annually

Significant unmet need in 1L treatment with current standard of care, ruxolitinib:

- <50% of patients achieve SVR35² and TSS50²
- ~25% of male patients achieve an SVR35 at week 24³
- ~23% patients who start on ruxolitinib 15 mg BID achieve an SVR35 at week 24⁴
 - ~70% patients in the real world start on ruxolitinib 15 mg BID or less⁵

Leading causes of ruxolitinib discontinuation are thrombocytopenia and anemia, which is associated with shorter survival^{6,7,8}


Selinexor has the Potential to Shift the Treatment Paradigm*

- ✓ XPO1 inhibition is a novel and potentially fundamental mechanism in MF
- ✓ Synergism with ruxolitinib observed in preclinical data⁹
- ✓ Rapid, deep and sustained spleen response, and robust symptom improvement across all subgroups
- ✓ Generally tolerable and manageable side effect profile enabling sustained therapy
- ✓ Disease modifying with rapid normalization of platelets and maintenance of hemoglobin levels


* Based on selinexor+ruxolitinib Ph 1 results using data cut as of February 24, 2023

The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Selinexor Can Inhibit Multiple Targets of the JAK/STAT Pathway, Enabling Independent Suppression of MF cells and Potentially Complementing the Function of JAKi's^{1,2,3,4,5}



Phase 1 Study (XPORT-MF-034¹) Evaluating Selinexor in Combination with Ruxolitinib in Treatment-naïve Myelofibrosis



* Enrollment completed; 24 patients had been assigned to either a 40 mg (n=10) or 60 mg (n=14) once weekly dose of selinexor, in combination with ruxolitinib 15/20 mg BID (twice daily)

Rapid, Deep and Sustained SVR35 and TSS50 Achieved with Selinexor 60 mg

Population	Timepoint	SVR35		TSS50	
		Selinexor 40 mg +ruxolitinib n (%)	Selinexor 60 mg +ruxolitinib n (%)	Selinexor 40 mg +ruxolitinib n (%)	Selinexor 60 mg +ruxolitinib n (%)
Efficacy Evaluable	Week 12	3/10 (30.0)	10/12 ⁱ (83.3)	6/9 ⁱⁱⁱ (66.7)	8/10 ^v (80.0)
	Week 24	4/8ⁱⁱ (50.0)	11/12 (91.7)	4/7^{iv} (57.1)	7/9^{vi} (77.8)
Intent-to- Treat	Week 12	3/10 (30.0)	10/14 (71.4)	6/10 (60.0)	8/12 (66.7)
	Week 24	4/10 (40.0)	11/14 (78.6)	4/10 (40.0)	7/12 (58.3)

ⁱ Two patients discontinued prior to Week 24.

ⁱⁱ One patient discontinued prior to week 12; one patient with missing data at week 12 who subsequently discontinued prior to week 24.

ⁱⁱⁱ One patient with missing data

^{iv} Two patients discontinued prior to Week 24 and one had missing data.

^v One patient discontinued prior to week 12; one patient with missing data at week 12, who subsequently discontinued prior to week 24.


^{vi} Two patients discontinued prior to Week 24 and one had missing data.

SVR35, spleen reduction volume ≥35%

TSS50, total symptom score ≥ 50. Note: Median TSS was calculated for each cycle, regardless of number of scores collected per cycle

The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.


All Evaluable Patients* Treated with Selinexor 60 mg Achieved an SVR35 at Anytime



* n=12; one patient discontinued prior to week 12; one patient with missing data at week 12 who subsequently discontinued prior to week 24.

The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Selinexor + Ruxolitinib Treatment Effective for All JAKi Naïve Patients, Across All Subgroups Including Males and Patients Who Started at Low Ruxolitinib Doses




The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Efficacy with Selinexor in Combination with Suboptimal Dose of Ruxolitinib (5 mg) Further Supports XPO1 as a Fundamental MoA in MF

Retrospective, Exploratory Analysis from Phase 1 Selinexor + Ruxolitinib Study (034)

Spleen Volume Reduction



Total Symptom Score Reduction¹



“Based on limited clinical data, long-term maintenance at a 5 mg twice daily dose has not shown responses and continued use at this dose should be limited to patients in whom the benefits outweigh the potential risks.” Jakafi (ruxolitinib) U.S. Package Insert, January 2023

The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

1. One patient with missing TSS50 score
2. Assigned selinexor starting dose
Data cut February 24, 2023


Generally Tolerable and Manageable Side Effect Profile

Treatment Emergent Adverse Events	Selinexor 40mg + ruxolitinib (N=10)	Selinexor 60mg + ruxolitinib (N=14)
Any grade, >25% overall		
Nausea	7 (70.0)	11 (78.6)
Anemia	4 (40.0)	9 (64.3)
Fatigue	6 (60.0)	8 (57.1)
Thrombocytopenia	4 (40.0)	9 (64.3)
Constipation	2 (20.0)	7 (50.0)
Headache	4 (40.0)	5 (35.7)
Vomiting	2 (20.0)	7 (50.0)
Neutropenia	2 (20.0)	5 (35.7)
Dyspnea	2 (20.0)	5 (35.7)
Decreased appetite	2 (20.0)	4 (28.6)
Dysgeusia	2 (20.0)	4 (28.6)
Hyponatremia	1 (10.0)	5 (35.7)
Grade 3+, >5% overall		
Anemia	3 (30.0)	6 (42.9)
Thrombocytopenia	1 (10.0)	4 (28.6)
Neutropenia	2 (20.0)	1 (7.1)
Atrial fibrillation	2 (20.0)	1 (7.1)
Back pain	0	2 (14.3)
Treatment-related adverse events leading to treatment discontinuations		
Thrombocytopenia, Grade 3	0	1 (7.1)
Peripheral Neuropathy	0	1 (7.1)

- Treatment related discontinuations due to cytopenias were low (n=1)
- 75% of nausea events were Grade 1
 - One patient experienced Grade 3 nausea (no antiemetic prophylaxis)
- In the 60mg cohort, 64% of patients received one prophylactic antiemetic
 - Amongst the subgroup who received one prophylactic antiemetic, 67% of patients experienced nausea (Grade 1 only) compared to 100% of those who did not receive prophylactic antiemetics (Grades 1-3)


The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Patients Experienced Improved Weight with Selinexor in Combination with Ruxolitinib



- Patients' median weight increase at Week 24 was 3 kg in the 40mg cohort and 2.5 kg in the 60mg cohort
- Despite nausea and vomiting incidence, patients generally did not experience weight loss

Potential Disease Modification With Normalization of Platelets and Restoration of Hemoglobin Levels




The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

Selinexor 60 mg in Combination with Ruxolitinib Has the Potential to Transform 1L Myelofibrosis Treatment Paradigms

- Selinexor 40 mg and 60 mg dose levels generally well tolerated and manageable allowing most patients to remain on therapy (up to 68 weeks as of data cutoff)
 - Only two treatment related discontinuations observed: Peripheral neuropathy (n=1) and thrombocytopenia (n=1)
- Rapid, deep, and sustained spleen response and robust symptom improvement in patients treated with 60 mg selinexor: SVR35: 78.6% ITT (91.7%, EE) and TSS50: 58.3% ITT (77.8%, EE) at week 24
 - SVR35: Observed in 100% of evaluable patients at anytime; rates consistent by gender and ruxolitinib starting dose
 - SVR35 and TSS50 observed amongst patients treated with suboptimal doses of ruxolitinib 5 mg
- Disease modification observed as evidenced by rapid normalization of platelets and restoration of hemoglobin levels
- Efficacy and safety data support 60 mg dose of selinexor as the recommended dose for Phase 3 study in combination with ruxolitinib

XPORT-MF-034 Phase 3 Design*



Randomization stratified by:

- Dynamic International Prognostic Scoring System (DIPSS) risk category intermediate -1 vs. intermediate -2 or high-risk
- Spleen volume <1800 cm³ vs. >1800 cm³ by MRI/CT scan
- Baseline platelet counts 100-200 x 10⁹/L vs. >200 x 10⁹/L

The safety and efficacy of selinexor in myelofibrosis has not been established and has not been approved by the U.S. FDA or any other regulatory authority.

MYELOYDYSPLASTIC NEOPLASMS



Eltanexor Has the Potential to Improve Survival in Higher Risk Relapsed/Refractory Myelodysplastic Neoplasms (MDS)

What are Myelodysplastic Neoplasms (MDS)?

- Blood-forming cells in marrow become abnormal and create immature blood cells that are not able to function properly


Treatment Landscape

- Hypomethylating agents (HMA) are the current standard of care (SoC) for patients with newly diagnosed, higher-risk MDS
- ~50% of 1L treated patients do not respond; for patients who achieve a response, responses last < 2 years²

Opportunity and Unmet Need

- No SoC exists for higher risk, relapsed or refractory MDS patients
- ~80% of patients treated in the community do not receive a 2L treatment
- Prognosis of higher risk relapsed/refractory disease is poor, with a median overall survival of four to six months^{3,4}
- Transfusion dependence is a major problem in R/R MDS patients; transfusions impair quality of life and cause potential complications
- Critical need for novel and more effective treatment options for this patient population

~12,000 - 20,000 patients diagnosed with higher-risk MDS each year in the US¹




1. What is MDS?, MDS Foundation, accessed April 21, 2021, <https://www.mds-foundation.org/what-is-mds/>

2. Gil-Perez A. Ther Adv Hematol. 2019 doi:10.1177/2040620719847059.

3. Jabbour E, Cancer. 2010;116(16):3830-4

4. Prébet T. J Clin Oncol. 2011;29:3322-7

Phase 2 Study of Eltanexor in Higher Risk, Relapsed/Refractory MDS




Encouraging Efficacy Observed in Higher Risk Relapsed/Refractory MDS; a Hard-to-Treat Patient Population with Limited Treatment Options

Data from Interim Analysis (n=30)	
	Eltanexor (ITT)
Overall Response Rate* (ORR = mCR + HI), n (%)	8/30 (26.7)
Marrow Complete Response (mCR)	8/30 (26.7)
Hematologic Improvement ($\pm$ mCR)	2/30 (6.7)
Disease Control Rate, n (%)	21/30 (70.0)
Transfusion Independence for Red Blood Cells and/or Platelets, n (%)	8/28** (28.6)

CI, confidence interval; HI, hematologic improvement; mCR, marrow complete response; NR, not reached; ORR, overall response rate as reported by Investigators

* In efficacy evaluable population: ORR: 31% and DCR: 81%

** 28 patients were RBC or platelet transfusion dependent at baseline



Prognosis of higher risk relapsed/refractory disease is poor, with a median overall survival of **four to six months**^{1,2}

Eltanexor is an investigational drug not approved by any regulatory authority; its safety and efficacy have not been established.

Generally Tolerable and Manageable Side Effect Profile

Treatment Emergent Adverse Events	Eltanexor (n=30)
Any grades, ≥20% overall	
Asthenia	14 (46.7)
Diarrhea	13 (43.3)
Nausea	10 (33.3)
Constipation	9 (30.0)
Neutropenia	9 (30.0)
Thrombocytopenia	8 (26.7)
Decreased appetite	7 (23.3)
Weight decreased	7 (23.3)
Contusion	6 (20.0)
Epistaxis	6 (20.0)
Oedema peripheral	6 (20.0)

Treatment Emergent Adverse Events	Eltanexor (n=30)
Grade 3+, ≥10%	
Neutropenia	9 (30.0)
Thrombocytopenia	8 (26.7)
Asthenia	5 (16.7)
Anemia	4 (13.3)
Febrile neutropenia	4 (13.3)
Leukopenia	3 (10.0)
Epistaxis	3 (10.0)
Fall	3 (10.0)

- There were no treatment-related AEs leading to death
- There were 3 patients that discontinued treatment due to a treatment-related AE: alanine aminotransferase (ALT) increase, aspartate aminotransferase (AST) increase, asthenia, and hemorrhagic diarrhea (increases in ALT and AST were experienced in one patient)

Eltanexor is an investigational drug not approved by any regulatory authority; its safety and efficacy have not been established.

PARTNERSHIPS, FINANCIAL HIGHLIGHTS AND MILESTONES



Strategic Partnerships Driving Expansion of Our Global Footprint

Commercial Partnerships Serving Key Global Markets



AsiaPac



EU, LatAm, Middle East & Africa



Israel



Canada

1Q 2023 Financial Results

Statements of Operations (\$ millions)	1Q 2023	1Q 2022
Total Revenue	\$38.7	\$47.7
U.S. XPOVIO Net Product Revenue	28.3	28.3
License and Other Revenue	10.4	19.4
Total Operating Expenses	\$69.6	\$82.3
Cost of Sales	1.4	1.4
Research and Development Expenses	32.3	42.1
Selling, General & Administrative Expenses	35.9	38.8
Net Loss	(\$34.1)	(\$41.4)
Net Loss per share	(\$0.30)	(\$0.53)

Balance Sheet (\$ millions)	March 31, 2023	Dec 31, 2022
Cash, Cash Equivalents Restricted Cash and Investments	\$261.9	\$279.7

Revised 2023 Financial Guidance

- Total Revenue of \$145-\$160M
- U.S. XPOVIO Net Product Revenue of \$110-\$125M
- Guidance revision primarily due to:
 - Significant increase in utilization of PAP (free drug) due to recent lack of funding to foundations for multiple myeloma patients
 - In 2024, significantly fewer patients expected to utilize PAP for co-pay assistance due to IRA related re-design of Part-D benefits
- Non-GAAP R&D and SG&A Expenses of \$245-\$260M¹
- Cash runway expected to be sufficient to fund planned operations into late 2025

Upcoming Milestones for 2023 and Beyond

MULTIPLE MYELOMA

- Leverage commercial capabilities and grow XPOVIO (2023)
- Continuation of global launches (2023)
- Report top-line results from pivotal Phase 3 study evaluating SPd¹ (2H 2024)
- Present data on XPOVIO pre/post T-cell therapy (2H 2023-1H 2024)

ENDOMETRIAL CANCER

- Present exploratory updated results from the *TP53* subgroup from the SIENDO study at a medical conference (2023)
- Report top-line results from pivotal Phase 3 study EC-042 in *TP53* wild-type EC (2H 2024)

MYELOFIBROSIS

- Report updated results in Phase 1 trial of selinexor+ ruxolitinib in treatment-naïve MF (1H 2023) ✓
- Initiate pivotal Phase 3 selinexor + ruxolitinib study in treatment-naïve MF (1H 2023)
- Define optimal mono and innovative combo development plan (1H 2023)

MYELODYSPLASTIC NEOPLASMS

- Report interim Phase 2 eltanexor data in higher risk R/R MDS (1H 2023) ✓
- Define development plan in higher risk R/R MDS (2H 2023)

1. Selinexor + pomalidomide + dexamethasone



Thank you!
