



A Commercial-Stage Pharmaceutical Company Pioneering Novel Cancer Therapies

February 2021

Forward-looking Statements and Other Important Information

This presentation contains forward-looking statements within the meaning of The Private Securities Litigation Reform Act of 1995. Such forward-looking statements include those regarding Karyopharm's guidance on its 2021 non-GAAP research and development and selling, general and administrative expenses; expectations and plans relating to XPOVIO for the treatment of patients with relapsed or refractory multiple myeloma or relapsed or refractory diffuse large B-cell lymphoma; commercialization of XPOVIO or any of its drug candidates and the commercial performance of XPOVIO; submissions to, and the review and potential approval of selinexor by, regulatory authorities, including the Company's regulatory strategy, the anticipated availability of data to support such submissions, timing of such submissions and actions by regulatory authorities and the potential availability of accelerated approval pathways; the expected design of the Company's clinical trials; and the therapeutic potential of and potential clinical development plans for Karyopharm's drug candidates, especially selinexor. Such statements are subject to numerous important factors, risks and uncertainties, many of which are beyond Karyopharm's control, that may cause actual events or results to differ materially from Karyopharm's current expectations. For example, there can be no guarantee that Karyopharm will successfully commercialize XPOVIO; that regulators will agree that selinexor qualifies for conditional approval in the E.U. as a result of data from the STORM study or confirmatory approval in the E.U. based on the BOSTON study in patients with multiple myeloma; or that any of Karyopharm's drug candidates, including selinexor, will successfully complete necessary clinical development phases or that development of any of Karyopharm's drug candidates will continue. Further, there can be no guarantee that any positive developments in the development or commercialization of Karyopharm's drug candidate portfolio will result in stock price appreciation. Management's expectations and, therefore, any forward-looking statements in this press release could also be affected by risks and uncertainties relating to a number of other factors, including the following: the risk that the COVID-19 pandemic could disrupt Karyopharm's business more severely than it currently anticipates, including by negatively impacting sales of XPOVIO, interrupting or delaying research and development efforts, impacting the ability to procure sufficient supply for the development and commercialization of selinexor or other product candidates, delaying ongoing or planned clinical trials, impeding the execution of business plans, planned regulatory milestones and timelines, or inconveniencing patients; the adoption of XPOVIO in the commercial marketplace; the timing and costs involved in commercializing XPOVIO or any of Karyopharm's drug candidates that receive regulatory approval; the ability to retain regulatory approval of XPOVIO or any of Karyopharm's drug candidates that receive regulatory approval; Karyopharm's results of clinical trials and preclinical studies, including subsequent analysis of existing data and new data received from ongoing and future studies; the content and timing of decisions made by the U.S. Food and Drug Administration and other regulatory authorities, investigational review boards at clinical trial sites and publication review bodies, including with respect to the need for additional clinical studies; the ability of Karyopharm or its third party collaborators or successors in interest to fully perform their respective obligations under the applicable agreement and the potential future financial implications of such agreement; Karyopharm's ability to obtain and maintain requisite regulatory approvals and to enroll patients in its clinical trials; unplanned cash requirements and expenditures; development or regulatory approval of drug candidates by Karyopharm's competitors for products or product candidates in which Karyopharm is currently commercializing or developing; and Karyopharm's ability to obtain, maintain and enforce patent and other intellectual property protection for any of its products or product candidates. These and other risks are described under the caption "Risk Factors" in Karyopharm's Quarterly Report on Form 10-Q for the quarter ended September 30, 2020, which was filed with the Securities and Exchange Commission (SEC) on November 2, 2020, and in other filings that Karyopharm may make with the SEC in the future. Any forward-looking statements contained in this presentation speak only as of the date hereof, and, except as required by law, Karyopharm expressly disclaims any obligation to update any forward-looking statements, whether as a result of new information, future events or otherwise. Karyopharm regularly uses its website to post information regarding its business, drug development programs and governance. Karyopharm encourages investors to use www.karyopharm.com, particularly the information in the section entitled "Investors," as a source of information about Karyopharm. References to www.karyopharm.com in this presentation are not intended to, nor shall they be deemed to, incorporate information on www.karyopharm.com into this presentation by reference. Other than the accelerated approval of XPOVIO, selinexor, eltanexor, KPT-9274 and verdinexor are investigational drugs that have not been approved by the FDA or any other regulatory agency, and the safety and efficacy of these drugs has not been established by any agency.

XPOVIO® and NEXPOVIO® are registered trademarks of Karyopharm Therapeutics Inc. Any other trademarks referred to in this presentation are the property of their respective owners. All rights reserved.

TABLE OF CONTENTS


- Overview of Karyopharm
- Overview of XPOVIO®
- Pipeline and Future Opportunities
- Additional Highlights and Next Steps
- Appendix: Clinical Data
- Appendix: Commercial Insights

Karyopharm at a Glance



Commercial-stage, global pharmaceutical company with **one FDA-approved drug in three oncology indications** and **three** additional drug candidates in clinical development

Industry leader in targeting **nuclear export dysregulation** as a mechanism to treat cancer




XPOVIO® (selinexor) received first accelerated approval from the FDA in July 2019 (penta-refractory multiple myeloma)

In December 2020, XPOVIO received expanded FDA approval in patients with multiple myeloma **as early as first relapse**



Karyopharm at a Glance




XPOVIO also received accelerated approval in June 2020 for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL)

Numerous key milestones expected over the next 12 months

All programs developed in-house

Ongoing clinical development for XPOVIO and next-generation programs in earlier lines of treatment, in combination trials, and in **additional tumor types** across both **hematologic and solid tumor malignancies**



XPOVIO Now Approved In Significantly Expanded Multiple Myeloma (MM) Patient Population



NEW INDICATION

XPOVIO[®]
(selinexor)

Now approved for adult patients with multiple myeloma as early as first relapse

XPOVIO is indicated in combination with bortezomib and dexamethasone for the treatment of adult patients with multiple myeloma who have received at least one prior therapy¹

FDA Expanded Approval on December 18, 2020

XPOVIO is the **FIRST and **ONLY** nuclear export / XPO1 inhibitor approved by the FDA**

Full Prescribing Information and Medication Guide available at www.XPOVIO.com

XPOVIO Received Accelerated Approval in Relapsed or Refractory DLBCL

XPOVIO is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL), not otherwise specified, including DLBCL arising from follicular lymphoma, after at least 2 lines of systemic therapy¹

This indication is approved under accelerated approval based on response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).



Full Prescribing Information and Medication Guide
are available at [XPOVIO.com](https://www.xpovio.com)

Safety Highlights from the XPOVIO Prescribing Information¹

No Black Box Warnings

No Contraindications

Patient Medication Guide

Warnings and Precautions

- Thrombocytopenia
- Neutropenia
- Gastrointestinal Toxicity
- Hyponatremia
- Serious Infection
- Neurological Toxicity
- Embryo-Fetal Toxicity
- Cataract

Monitoring Instructions and Recommended Concomitant Treatments

- Monitor complete blood count (CBC) with differential, standard blood chemistries, body weight, nutritional status, and volume status at baseline and during treatment as clinically indicated. Monitor more frequently during the first three months of treatment.
- Patients advised to maintain adequate fluid and caloric intake throughout treatment. Consider intravenous hydration for patients at risk of dehydration.
- Provide prophylactic antiemetics. Administer a 5-HT₃ receptor antagonist and other anti-nausea agents prior to and during treatment with XPOVIO
- Recommended XPOVIO dosage reductions and dosage modifications for adverse reactions are included in the Prescribing Information

Q4 2020 and Recent Achievements Help Position Karyopharm For Significant Future Growth



Commercial Update

- **Expanded U.S. approval** of XPOVIO® and commercial launch (Dec 18, 2020)
- Q4 and full year 2020 XPOVIO net sales of **\$20.2M** and **\$76.2M**, respectively
- National Comprehensive Cancer Network® adds **three XPOVIO treatment regimens** to its clinical practice guidelines for multiple myeloma
- **>170** and **>700 new** physicians / accounts prescribed XPOVIO in Q4 and full year 2020, respectively



Pipeline / Clinical Data Update


- **Positive CHMP opinion** issued for NEXPOVIO® (selinexor) MAA in Europe based on STORM study (multiple myeloma)
- **Regulatory approval** in Israel
- **Positive Phase 3 BOSTON study** results published in the *Lancet* (multiple myeloma)
- **Positive Phase 3 Seal study** results presented at CTOS annual meeting (liposarcoma)
- **Phase 3 SIENDO study** passes planned interim futility analysis (endometrial cancer)
- **1st patient dosed** in Phase 3 confirmatory trial in DLBCL (XPORT-DLBCL-30)



Corporate Development and Balance Sheet

- Q4 and full year 2020 license revenue of **\$14.9M** and **\$31.9M**, respectively
- Ended Q4 2020 with **\$276.7M** in cash, cash equivalents, restricted cash and investments
- Cash runway expected to be sufficient to fund planned operations into **late 2022**

XPOVIO 2020 Quarterly Net Product Sales and Prescription (RX) Trends



Update on Q4 2020 and Recent Sales Trends

- Q4 XPOVIO sales primarily affected by:
 1. Surge in U.S. COVID-19 cases
 - Fewer patient visits to their health care providers
 - Reduced ability of Karyopharm commercial team to access customers in-person
 2. Increased competition in the penta-refractory multiple myeloma and 3rd line+ DLBCL patient settings
- Strong return to sales growth seen in December and January following declines in October and November of 2020
 - Positive sales momentum seen in December 2020 and January 2021 following NCCN guideline update adding three XPOVIO regimens to its treatment recommendations and the expanded XPOVIO FDA approval granted on December 18th
 - Increase in estimated new patient starts seen in December and January as compared to October and November

Status Update on Commercial Expansion in Europe

1. Positive CHMP opinion in Europe issued on January 29, 2021

- Positive opinion highlights the CHMP's recognition of the positive clinical benefit-risk profile for oral NEXPOVIO (expected brand name for selinexor in Europe)
- European Commission's final decision on the NEXPOVIO MAA expected by April 2021
- Karyopharm to submit second regulatory filing based on data from the confirmatory Phase 3 BOSTON study by April 2021

2. Selinexor added to European treatment guidelines in November 2020

- Multiple myeloma treatment guidelines developed by the European Hematology Association (EHA) and European Society for Medical Oncology (ESMO) added SVd* as a treatment option in the second line setting and Sd** for patients with triple-class refractory disease

3. Advancing partnership discussions for international markets


- Evaluating potential collaboration strategies with commercial partners for Europe and Japan
- Largest revenue opportunity in Europe to come following potential approval based on data from the BOSTON study which could occur before the end of 2021¹

*Selinexor, Velcade®, dexamethasone

** Selinexor, dexamethasone

INNOVATIVE
APPROACH TO
TARGETING
CANCER

Core Pillars of Cancer Drug Therapy




XPOVIO® (selinexor) / SINE Mechanism of Action: Inhibition of XPO1¹⁻⁴

XPO1 OVEREXPRESSION

- Enables cancer cells to escape tumor suppressor proteins (TSPs) mediated cell cycle arrest and induction of apoptosis
- Correlates with poor prognosis and drug resistance

INHIBITION OF XPO1 IMPACTS TUMOR CELLS VIA 3 CORE MECHANISMS

1. Increases nuclear levels and activation of TSPs
2. Traps oncoprotein mRNA in the nucleus leading to reduced oncoprotein levels
3. Retains activated glucocorticoid receptor in the nucleus



MULTIPLE MYELOMA




XPOVIO's December 2020 Approval (BOSTON Study) Expands Addressable Population by 5X

2020 U.S. MM Epidemiology¹

~130K total prevalence

~60,000 patients not on treatment / in long-term remission



BOSTON approval:
33,000 (in the 2nd and 3rd Line)



Expands addressable population by 5X

STORM approval:
~6,000 (in the 4th Line+)



¹ Sources: Decision Resources used for growth rates for treated patients and overall prevalence; Kantar Cancer Impact; SEER; Cancer Facts and Figures 2020

XPOVIO's December 2020 Approval (BOSTON Study) Extends Duration of Treatment by **3X**

	Median # of Prior Therapies	XPOVIO Dose Frequency	ORR	Median Progression Free Survival (in months)	Mean Duration of Treatment (in months)
STORM Study^{1,2} (Penta-Refractory)	8	Twice per week (in combination with dexamethasone)	25%	3.7	3
BOSTON Study³ (1–3 Prior Therapies)	2	Once per week (in combination with once-weekly Velcade and dexamethasone)	76%	13.9	10

BOSTON approval extends duration of treatment by **3X** 


¹ STORM study provided the basis for XPOVIO's approved indication in combination with dexamethasone for the treatment of adult patients with relapsed or refractory multiple myeloma (RRMM) who have received at least 4 prior therapies and whose disease is refractory to at least 2 proteasome inhibitors (PI), at least 2 immunomodulatory agents (IMiD), and an anti-CD38 monoclonal antibody (mAb). ² XPOVIO Prescribing Information and Chari et al., NEJM. August 2019. ³ Grosicki S, et al. Lancet. 2020.

Progression Free Survival (PFS) Significantly Longer with XVd Compared to Vd

30% reduction in risk of progression or death¹

Hazard ratio: 0.70
(96% CI 0.5-0.93), $p=0.0075$

ONCE-WEEKLY, ORAL XPOVIO + VD DELIVERED AN EARLY AND SUSTAINED PFS ADVANTAGE VERSUS TWICE-WEEKLY Vd¹



Hazard ratio (HR) is based on stratified Cox's proportional hazard regression modeling, p-value based on stratified log-rank test.

*According to the International Myeloma Working Group (IMG Uniform Response Criteria for Multiple Myeloma, as assessed by an Independent Review Committee (IRC). XVd=XPOVIO® (selinexor) with Velcade® (bortezomib) and dexamethasone; Vd=Velcade and dexamethasone.

Overall Response Rates (ORR) Demonstrated in the BOSTON Trial¹:

Responses observed with oral, once-weekly XPOVIO + Vd were rapid and durable versus twice-weekly Vd¹

MEDIAN
TIME TO
RESPONSE¹

1.4

MONTHS
once-weekly
XPOVIO + Vd

VS

1.6

MONTHS
twice-weekly
Vd

MEDIAN
DURATION OF
RESPONSE¹

20.3

MONTHS
once-weekly
XPOVIO + Vd

VS

12.9

MONTHS
twice-weekly
Vd

Depth of response observed with once-weekly XPOVIO + Vd was significant versus twice-weekly Vd (p=0.0082)¹

ONCE-WEEKLY XPOVIO + VD

≥VGPR: 44.6%

PR:32%

VGPR:28%

sCR+CR: 17%

ORR: 76.4%

TWICE-WEEKLY VD

≥VGPR: 32.4%


PR:30%

VGPR:22%

sCR+CR: 10%

ORR: 62.3%

IMPROVEMENT IN
ORR WAS OBSERVED
ACROSS A VARIETY OF
PATIENT SUBGROUPS²



Adverse Reactions in Patients with Multiple Myeloma Who Receive XVd¹

- The most common adverse reactions ($\geq 20\%$) in patients with multiple myeloma who receive XVd are fatigue, nausea, decreased appetite, diarrhea, peripheral neuropathy, upper respiratory tract infection, decreased weight, cataract and vomiting
- Grade 3-4 laboratory abnormalities ($\geq 10\%$) are thrombocytopenia, lymphopenia, hypophosphatemia, anemia, hyponatremia and neutropenia

Full Prescribing Information and Medication Guide available at www.XPOVIO.com

Key Messaging for Expanded Indication in Multiple Myeloma

Unmet Need

In MM, treating with *different mechanisms as early as possible* may be *vital for success*

Efficacy & Duration

Weekly XPOVIO + Vd conferred a rapid and sustained PFS benefit. And patients achieved a clinically significant durable response with once-weekly XPOVIO + Vd regardless of cytogenetics, renal impairment, or prior therapeutic exposure

Mechanism of Action

XPOVIO is the first and only FDA-approved oral XPO1 inhibitor that gets to the cell's nucleus which leads to cell cycle arrest and apoptosis in cancer cells

- First new mechanism approved since 2016 for the treatment of MM in patients who received at least 1 prior therapy
- XPOVIO has a strong synergistic effect with proteasome inhibitors, leading to cancer cell death

Dosing


The oral, once-weekly XPOVIO + once-weekly Vd combination offers a high-efficacy regimen while potentially reducing the burden for in-office Velcade® treatment

Safety

XPOVIO + Vd offers a manageable safety profile for a broad range of patients

Specific Patient Types that Physicians May Find Particularly Appealing for the XPOVIO + Vd Regimen in the 2nd Line+ Settings

- Received Revlimid® and Darzalex® in the front-line setting and are Velcade®-naïve following first relapse
- Received only a short course of Velcade® in the front-line setting prior to stem cell transplant
- Have high-risk disease and/or cytogenetic abnormalities
- Have renal dysfunction
- Prefer a once weekly oral drug and once-weekly injection rather than IV infusions or more frequent visits to the clinic
- Might benefit from a drug with a completely novel mechanism of action, synergistic with a proteasome inhibitor





DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL)

Commercial Strategy for XPOVIO in DLBCL

XPOVIO POSITIONING

- Position XPOVIO as the preferred DLBCL treatment option after two prior lines of therapy instead of traditional intravenous chemotherapy by educating physicians on the deep and durable efficacy achieved in clinical studies with oral, single-agent, novel, XPOVIO
- XPOVIO offers compelling efficacy with a manageable safety profile and is now:
 - **First** oral therapy approved for RR DLBCL
 - **First** single agent approved in any line of DLBCL treatment
 - **First** therapy a RR DLBCL patient can take at home

Key Features of XPOVIO for the Treatment of Patients With RR DLBCL

Factors that influence treatment choice	Key features of XPOVIO
• Clinical efficacy	• 29% ORR¹ • 13% CR¹ • Clinically meaningful duration of response¹
• Previous therapies / approaches	• Novel mechanism of action
• Subtype and histology	• Similar efficacy seen across both ABC and GCB patient sub-types
• Comorbidities • Functional status, age, frailty	Common adverse events do not include: • Peripheral neuropathy • Cardiac, liver or kidney toxicity • Opportunistic infections
• Patient preferences / logistical dynamics	• Oral route of administration taken only twice per week • Single agent, not combined with chemotherapy

PIPELINE
AND SOLID
TUMORS

Karyopharm's Novel Pipeline | Selinexor

HEMATOLOGIC MALIGNANCIES STUDY NAME	PHASE 1	PHASE 2	PHASE 3
Multiple Myeloma (relapsed/refractory) STORM	APPROVED ¹		
Multiple Myeloma (relapsed/refractory) BOSTON ²	APPROVED ¹		
Diffuse Large B-cell Lymphoma (relapsed/refractory) SADAL	APPROVED ¹		
Multiple Myeloma (relapsed/refractory and front-line) STOMP ³			
Diffuse Large B-cell Lymphoma (combination with rituximab-gemcitabine-dexamethasone-platinum (R-GDP)) XPORT-DLBCL-030 (Phase 2/3)			
Diffuse Large B-cell Lymphoma (combination with chemo and non-chemo regimens) XPORT-DLBCL-025 ⁴			
Myelofibrosis (previously treated) XPORT-MF-035 ⁴			
Myelofibrosis (combination with ruxolitinib) XPORT-MF-034 ⁴			

Additional new Phase 3 study (XPORT-MM-031) evaluating XPOVIO in combination with pomalidomide and dexamethasone in patients with previously treated multiple myeloma expected to start in **2021**

Karyopharm's Novel Pipeline | Selinexor

SOLID TUMOR MALIGNANCIES STUDY NAME	PHASE 1	PHASE 2	PHASE 3
Liposarcoma (advanced unresectable dedifferentiated liposarcoma) SEAL			
Endometrial Cancer (maintenance therapy) SIENDO			
NSCLC (combination with docetaxel) XPORT-STP-027			
CRC (combination with pembrolizumab) XPORT-STP-027			
GLIOBLASTOMA MULTIFORME (GBM) STUDY NAME	PHASE 1	PHASE 2	PHASE 3
Glioblastoma (recurrent gliomas) KING			
Glioblastoma (combination with active agents / newly diagnosed or recurrent) XPORT-GBM-029			

Additional new Phase 2 study evaluating XPOVIO in combination with pembrolizumab in patients with newly diagnosed or recurrent metastatic melanoma expected to start in **2021**

Karyopharm's Novel Pipeline | Eltanexor, KPT-9274 and Verdinexor

ADDITIONAL ONCOLOGY PROGRAMS - ELTANEXOR AND KPT-9274	PHASE 1	PHASE 2	PHASE 3
Myelodysplastic Syndromes (MDS) (single agent or in combination with hypomethylating agents) ¹ Drug: Eltanexor			
Colorectal Cancer (CRC) and Prostate Cancer (PrC) Drug: Eltanexor			
Solid Tumors & AML Drug: KPT-9274			
INFECTIOUS DISEASES & AUTOIMMUNE DISORDERS - VERDINEXOR	PHASE 1	PHASE 2	PHASE 3
Healthy Human Volunteers			
Systemic Lupus Erythematosus (SLE) VALOR-SLE-705 ²			
Severe Influenza, Respiratory Syncytial Virus (RSV), HIV Inflammation, Spinal Cord Injury ²			

Solid Tumor Update: SEAL Phase 3 Positive Top-Line Results

TOP-LINE PHASE 3 DATA

- Study met **primary endpoint** with significant increase in progression-free survival in patients with unresectable dedifferentiated liposarcoma following at least two prior therapies
- Hazard ratio=**0.70**; p=**0.023**
- Safety profile consistent with previous clinical studies with fewer hematologic and infectious adverse events as compared to selinexor studies in patients with multiple myeloma and diffuse large B-cell lymphoma
- Full data presented in an oral presentation at the Connective Tissue Oncology Society (CTOS) Annual Meeting on November 20, 2020

STRATEGIC IMPLICATIONS

- Positive pivotal data in liposarcoma demonstrates XPOVIO's substantial potential across multiple solid tumors, representing a major advance for the development and commercial potential of XPOVIO in oncology
- Consistent with other, earlier stage positive results from ongoing XPOVIO studies in diseases such as endometrial cancer, GBM, melanoma, lung cancer, and others

ADDITIONAL HIGHLIGHTS AND NEXT STEPS



Current Partnerships

Commercial partnerships to serve global markets

Antengene Corporation

Licensing partner for selinexor, eltanexor, verdinexor and KPT-9274 in China, South Korea, Taiwan, Australia and other Asia-Pacific markets, with the exception of Japan

Neopharm Group

Exclusive distribution agreement for the commercialization of XPOVIO in Israel and the Palestinian Authority

FORUS Therapeutics, Inc.

Exclusive distribution agreement for the commercialization of XPOVIO in Canada

Europe, Japan and Other Key Markets

Evaluating potential collaboration arrangements with commercial partners; analyzing potential for Karyopharm to commercialize in select European markets

KARYOPHARM IS COMMITTED TO WORKING ACROSS THE GLOBE TO BRING NOVEL THERAPIES TO PATIENTS

Q4 and Full Year 2020 Financial Results

Statement of Operations	Three Months Ended December 31 st		Twelve Months Ended December 31 st	
	2020	2019	2020	2019
Total Revenue	\$35.1M	\$18.1M	\$108.1M	\$40.9M
XPOVIO Net Sales	\$20.2M	\$17.7M	\$76.2M	\$30.5M
License and other Revenue	\$14.9M	\$0.4M	\$31.9M	\$10.4M
Total Operating Expenses	\$72.2M	\$61.4M	\$279.9M	\$230.2M
Cost of Sales	\$1.1M	\$1.4M	\$2.7M	\$2.4M
Research and Development Expenses	\$37.2M	\$31.6M	\$150.8M	\$122.3M
Selling, General & Administrative Expenses	\$33.9M	\$28.4M	\$126.4M	\$105.4M
Net Loss	\$43.4M (\$0.59 per share)	\$48.6M (\$0.76 per share)	\$196.3M (\$2.72 per share)	\$199.6M (\$3.22 per share)

Balance Sheet and Financial Guidance

Balance sheet	December 31, 2020	December 31, 2019
Cash, Cash Equivalents, Restricted Cash and Investments	\$276.7M	\$265.8M

- Non-GAAP R&D and SG&A expenses are expected to be in the range of \$280-300M for the full year 2021¹
 - Cash runway expected to be sufficient to fund planned operations into late 2022

¹ Excludes stock-based compensation expense. This outlook can only be provided on a non-GAAP basis because Karyopharm cannot reliably predict without unreasonable efforts the timing or amount of the factors that substantially contribute to the projection of stock compensation expense.

Numerous Expected Key Milestones in 2021

1H 2021

1. CHMP opinion on STORM MAA ✓
2. Start of confirmatory Phase 3 Study in DLBCL in support of 2020 accelerated approval ✓
3. EMA conditional approval based on STORM data¹
4. EMA submission of BOSTON study data (Type II variation)
5. Increased U.S. XPOVIO sales following expanded FDA approval in multiple myeloma

2H 2021

1. SIENDO Phase 3 study fully enrolled and topline data announced
2. EMA expanded approval based on BOSTON study¹
3. Initiation of Phase 3 study evaluating XPOVIO + pomalidomide in patients with multiple myeloma
4. Initiation of Phase 2 study evaluating XPOVIO + pembrolizumab in patients with metastatic melanoma
5. Additional combination data with XPOVIO and other standard of care anti-cancer drugs to be presented at key medical meetings
6. Continued, increased U.S. XPOVIO sales

A photograph of an elderly couple standing outdoors at sunset. The man, on the right, has white hair and is wearing a light blue button-down shirt over a grey t-shirt. The woman, on the left, has dark hair and is wearing a light pink ribbed top. They are both smiling and looking towards the right. The background is a bright, hazy sunset with trees and a building visible in the distance.

QUESTIONS?

APPENDIX:
CLINICAL DATA

Overview of Efficacy Data for Accelerated Approval of XPOVIO (n=83)¹

25.3% Overall Response Rate (ORR)

INCLUDING

1 Stringent complete response

0 Complete responses

4 Very good partial responses

16 Partial responses

Full STORM results from all 122 patients published in NEJM


- Median time to response: **4** weeks
- Median duration of response: **3.8** months



Additional Efficacy Data from Part 2 of STORM (n=122)¹


Change in M-Protein Levels²

71% of patients had a reduction in disease burden



Additional Efficacy Data from Part 2 of STORM (n=122)¹

Overall Survival by Group²



* ≥MR = Patients had a minor response or better; at least a 25% decrease in M protein


** PD/NE = Patients had progressive disease or disease not evaluable

Overview of Safety Data from STORM

- The most common adverse reactions ($\geq 20\%$) in patients with multiple myeloma who receive Xd (XPOVIO and dexamethasone) are thrombocytopenia, fatigue, nausea, anemia, decreased appetite, decreased weight, diarrhea, vomiting, hyponatremia, neutropenia, leukopenia, constipation, dyspnea and upper respiratory tract infection
- In the STORM trial, fatal adverse reactions occurred in 9% of patients. Serious adverse reactions occurred in 58% of patients. Treatment discontinuation rate due to adverse reactions was 27%.

Full Prescribing Information and Medication Guide are available at www.XPOVIO.com


BOSTON Study: Phase 3, Global, Randomized, Open Label, Controlled Study in Patients with Multiple Myeloma who Had Received 1-3 Prior Therapies




CR= complete response, DoR = duration of response, IMWG = International Myeloma Working Group, IRC = Independent Review Committee, OS = overall survival, PD = progressive disease, PFS = progression free survival, PR = partial response, PN = peripheral neuropathy, sCR = stringent complete response, TTNT = time to next therapy, VGPR = very good partial response. PFS defined as: Time from date of randomization until the first date of progressive disease, per IMWG response criteria, or death due to any cause, whichever occurred first, as assessed by IRC. ORR: Any response ≥PR (ie, PR, VGPR, CR, or sCR) based on the IRC's response outcome assessments, according to IMWG response criteria (Kumar et al. Lancet oncology 2016). All changes in MM disease assessments were based on baseline MM disease assessments.

*Vd weekly dosing and schedule for cycles≥ 9 as per SVd arm description.

Consistent PFS Benefit for XVd Across Subgroups



XVd Was Associated With a Significantly Higher ORR Overall and Across Patient Subgroups




One-sided P values for the Cochran-Mantel-Haenszel Test based on unstratified model. Data cut-off February 18, 2020.

ORR = Overall Response, based on Independent Review Committee's (IRC) response outcome assessments, according to IMWG response criteria (Kumar et al. Lancet Oncology 2016).

All changes in MM disease assessments were based on baseline MM disease assessments.


Overall Survival Interim Analysis (109 Deaths [27%])



SVd Arm	195	186	171	155	145	135	129	113	91	62	37	17	10	6	1	1	1	0
Vd Arm	207	193	185	169	156	149	141	125	96	62	41	19	11	7	0			

OS = Overall Survival, Data cut-off February 18, 2020.


Peripheral Neuropathy Rates were Significantly Lower with XVd than with Vd



STOMP: Study Overview & Objectives

Selinexor and backbone Treatments Of multiple Myeloma Patients

(STOMP): Multi-center, open-label, dose escalation (Phase 1) and expansion (Phase 2) study to assess the MTD, efficacy, and safety of selinexor in patients with RRMM



Additional XPOVIO Triplet Regimens Indicate Additive or Synergistic Activity Compared to Benchmark Doublet Regimens

Selinexor is currently being studied in the ongoing STOMP Phase 1b/2 trial evaluating selinexor and low-dose dexamethasone in combination with one of several standard approved myeloma therapies in patients with relapsed or refractory multiple myeloma

STOMP TRIAL			BENCHMARK DATA	
STOMP Triplet Regimen	# of Patients Treated to Date	Efficacy Data	Benchmark Regimen	Efficacy Data
Selinexor + Kyprolis + dex	24 (median 3 lines of prior therapy)	ORR = 75% ¹	Kyprolis + dex	ORR = 23% ⁵
Selinexor + Darzalex + dex	30 (Darzalex-naïve)	ORR = 73% ² PFS = 12.5 months ²	Darzalex	ORR = 29% ⁶ PFS = 3.7 months ⁶
Selinexor + Pomalyst + dex	60 (Pomalyst-naïve or non-refractory)	ORR = 54% (all pts) ³ ORR = 60% (pts dosed at RP2D) ³ PFS = 12.3 months ³	Pomalyst + dex	ORR = 29% ⁷ PFS = 3.6 months ⁷
Selinexor + Revlimid + dex	12 (Revlimid-naïve)	ORR = 92% ⁴	Revlimid + dex	ORR = 67% ⁸

Note: The STOMP study does not include any arms studying current myeloma backbone therapies without the combination of selinexor. The purpose of displaying the “Benchmark” data above is to highlight that the clinical results from STOMP reported to date strongly support ongoing / additional clinical investigation of selinexor in combination regimens.

In December 2020, the National Comprehensive Cancer Network® (NCCN) added three different XPOVIO combination regimens (Velcade®, Pomalyst® and Darzalex®) to its Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for previously treated multiple myeloma

¹ Gasparetto C, et al. ASH 2020. Abstract 1366. ² Gasparetto C, et al. ASCO 2020. Abstract 8510. ³ Chen C, et al. ASH 2020. Abstract 726. ⁴ White D, et al. ASH 2020. Abstract 1393. ⁵ Kyprolis Package Insert; Study PX-171-003 A1 ⁶ Lonial et al. Lancet 2016. ⁷ Pomalyst Package Insert. ⁸ Stewart et al. NEJM 2015. ⁹ Revlimid® (lenalidomide), Pomalyst® (pomalidomide), Velcade® (bortezomib), Kyprolis® (carfilzomib) or Darzalex® (daratumumab).

SADAL: A Phase 2b Study In DLBCL

Relapsed or Refractory or Transformed DLBCL

- Study included patients with at least two prior multi-agent therapies and who were ineligible for transplantation
- Included patients with Germinal Center B-Cell (GCB) and non-GCB subtypes

N=134

Oral Selinexor 60
mg

selinexor twice weekly
(4 week cycle)

SADAL: A Phase 2b Study In DLBCL

Selinexor 60mg twice weekly (n=134*)

OVERALL RESPONSE RATES^{1,2}

All Patients

29%

CRs 13%
PRs 16%

Genetic Subsets

34%

GCB (n=59)

21%

non-GCB (n=63)

ADDITIONAL ENDPOINTS²

Duration of Response

- Median DOR was **9.3** months
- Most responses at first scan (~**2** months)

Overall Survival

- All patients (n=127) = **9.1** months
- Patients with CR or PR (n=36) = **Not Yet Reached**
- Patients with stable disease (n=11) = **18.3** months
- Patients with Progressive Disease or No Response (n=80) = **4.3** months

Safety:

- Most common treatment related non-hematologic adverse events (AEs) were fatigue, nausea, decreased appetite, and diarrhea, primarily Grade 1/2, and most were manageable with dose modifications and/or supportive care
- Most common Grade 3/4 AEs were thrombocytopenia, lymphopenia, neutropenia, and anemia, and most were also manageable with dose modifications and/or supportive care

A photograph of a doctor in a white lab coat with a stethoscope around their neck, sitting at a desk with their hands clasped. A red semi-transparent box is overlaid on the right side of the image, containing the text 'APPENDIX: COMMERCIAL INSIGHTS'.

APPENDIX: COMMERCIAL INSIGHTS

Highly Experienced Team Educating the Market About XPOVIO

CUSTOMER-FACING FIELD FORCE

~70 sales representatives and nurse liaisons supporting commercial launch

- **~20** average years of pharmaceutical experience
- **~12** average years of hematology / oncology experience
- **~5** average years of MM experience

Experienced account management team responsible for payors and distribution partners

Extensive patient and HCP support program anchored by KaryForward™ platform

PRESCRIBER BASE¹

Multiple Myeloma


- **~400** accounts generate **~50%** of all prescriptions
- **~1,300** accounts generate **~80%** of all prescriptions

DLBCL

- **~3,000** physicians treat **~80%** of U.S. DLBCL patients
- **~75%** of targeted physicians are community-based
- **>50%** overlap between Karyopharm's multiple myeloma and DLBCL targets

Key XPOVIO Patient Metrics in 2020

Prescription Refill Rate for 1st and 2nd Prescription
(Only Includes Patients Eligible for a Refill)¹



2.9¹

Average Treatment
Cycles (RXs) Per Patient

13%¹

Patient Discontinuation
Rate Due to Side Effects

Drugs with Stand-Alone ($\pm$ Steroid) Anti-MM Activity and Approved for 1st or 2nd Line Treatment have Achieved >\$1B in Annual Sales

2020 Worldwide Sales¹

Immunomodulatory Agents

- **Revlimid[®]: \$12B**
- **Pomalyst[®]: \$3B**

Proteasome Inhibitors

- **Velcade[®]: >\$1B**
- **Kyprolis[®]: >\$1B**

Monoclonal Antibodies

- **Darzalex[®]: >\$4B**

Total U.S. Drug Sales in Relapsed or Refractory DLBCL Expected to Grow from \$762M in 2018 to Over \$3B by 2028

